

# Uremic toxins and peritoneal dialysis

NORBERT LAMEIRE, RAYMOND VANHOLDER, and RITA DE SMET

*Renal Division, Department of Medicine, University Hospital, Gent, Belgium*

**Uremic toxins and peritoneal dialysis.** Uremic toxicity is related in part to the accumulation of toxic substances, the nature of which has only partly been characterized. Because of the use of a highly permeable membrane and better preservation of the residual renal function, it could be anticipated that some of these uremic toxins are more efficiently cleared across the peritoneal membrane, and that the plasma and tissue levels of these compounds are lower than in hemodialysis patients. This article analyzes the generation and removal of several uremic toxins in peritoneal dialysis patients. The following uremic toxins are discussed:  $\beta_2$ -microglobulin, advanced glycation end products, advanced oxidation protein products, granulocyte inhibitory proteins, p-Cresol, and hyperhomocysteinemia. Some recent studies are reviewed suggesting that uremic toxins are involved in the progression of renal failure and are at least partially removed by peritoneal dialysis. We conclude that, although the plasma levels of some of these compounds are lower in peritoneal dialysis versus hemodialysis patients, it does not mean that the peritoneal dialysis patient is “better” protected against the numerous disturbances caused by these toxins.

It is now well accepted that the compounds accumulating in the uremic blood and tissues during the development of end-stage renal disease have either a direct or indirect impact on several biochemical and physiologic functions, as recently reviewed by Bergström [1] and our group [2, 3].

The recognition that uremic complications are related at least in part to the accumulation of these toxic substances has led to efforts not only to identify them or to inhibit their generation but also to enhance their detoxification or their removal by dialysis procedures. Much attention has been paid to the effect of hemodialysis on the removal of these toxins, while relatively less is known on their behavior in peritoneal dialysis.

However, because of the use of highly permeable peritoneal membranes and better preservation of the residual renal function in peritoneal dialysis, it could be anticipated that some of these uremic retention solutes are cleared more efficiently in peritoneal dialysis compared to hemodialysis.

**Key words:** end-stage renal disease, residual renal function, hyperhomocysteinemia, p-Cresol,  $\beta_2$ -microglobulin, advanced glycation end products, advanced oxidation protein product.

© 2001 by the International Society of Nephrology

This review will attempt to summarize the patterns of generation and removal of several uremic toxins in peritoneal dialysis patients.

For reasons of space, the discussion will be limited to  $\beta_2$ -microglobulin ( $\beta_2m$ ), advanced glycosylation end products (AGEs), advanced oxidation protein products (AOPPs), granulocyte inhibitory proteins, p-Cresol, and homocysteine. The potential role of the peritoneal elimination of some of these toxins on the progression of chronic renal failure, that is, the decline in residual renal function, will be discussed as well.

## $\beta_2$ -MICROGLOBULIN

Several studies have demonstrated lower plasma levels of  $\beta_2m$  in peritoneal dialysis patients than in hemodialysis patients and this has mainly been explained by a better preservation of the residual renal function in peritoneal dialysis patients [4–7]. An inverse relationship between blood levels of  $\beta_2m$  and residual renal function has been observed repeatedly in peritoneal dialysis [5, 6].

However, some studies found also lower blood levels for  $\beta_2m$  in peritoneal dialysis patients when patients were matched for residual renal function [7].

The most important biological alteration that has been related to  $\beta_2m$  is dialysis-related amyloidosis (DRA), which is characterized by amyloid deposition mainly in bone, joint and tendon structures, presenting as carpal tunnel syndrome, destructive arthropathy, and subchondral bone erosions and cysts.  $\beta_2$ -microglobulin has been demonstrated to be a major constituent of DRA-related amyloid fibrils. Recent studies have suggested a pathogenic role of a newly modified  $\beta_2m$ , that is, the advanced glycation end products (AGE) formed with carbonyl compounds derived from autooxidation of both carbohydrates and lipids (“carbonyl stress”) [reviewed in 8]. The lower blood levels of  $\beta_2m$  in peritoneal dialysis patients may explain the lower prevalence of  $\beta_2m$ -related amyloidosis. For example, Nomoto et al registered carpal tunnel syndrome in only 7 patients among a total of 5050 CAPD patients (0.14%) in a multicenter study in Japan [9]. In a recent multicenter European autopsy study, the incidence of amyloid in the peritoneal dialysis patients

was also lower compared to hemodialysis patients, even when the patients were matched for duration of dialysis therapy, although the difference was not statistically significant [10].

On the other hand, Tan et al performed serum amyloid component P scintigraphy, a specific technique for the detection of amyloid deposits, and found that the prevalence of positive scans in peritoneal dialysis patients was similar to that in hemodialysis patients [11].

The effects of 3-deoxyglucosone (3-DG), a potent protein cross-linking intermediate of the Maillard reaction, on the AGE modification of  $\beta_2m$  were studied, and the serum levels of 3-DG in patients undergoing hemodialysis, continuous ambulatory peritoneal dialysis (CAPD) and undialyzed patients were quantitated [12]. The serum levels of 3-DG were markedly increased in the dialyzed and undialyzed uremic patients. Incubation of  $\beta_2m$  with 3-DG at 37°C resulted in the emission of fluorescence characteristic for AGE, and caused AGE modification and dimer formation of  $\beta_2m$ , as demonstrated by Western blotting. The AGE-modified dimer of  $\beta_2m$  could be extracted from the amyloid tissue of a patient with DRA. 3-DG showed more intense and faster reactivity with  $\beta_2m$  to form AGE and the dimer as compared with glucose, and aminoguanidine suppressed the AGE and dimer formation of  $\beta_2m$  by 3-DG. It was concluded that the 3-DG accumulating in uremic serum may be involved in the AGE modification of  $\beta_2m$  amyloid.

## ADVANCED GLYCATION END PRODUCTS

Advanced glycation end products (AGEs) are formed during nonenzymatic glycation and oxidation (glycoxidation) reactions. AGE accumulation in uremia cannot be attributed to hyperglycemia nor simply to a decreased removal by glomerular filtration. Recent evidence has suggested that, in uremia, the increased carbonyl compounds derived from carbohydrates and lipids modify proteins not only by glycoxidation reaction, but also by a lipoxidation reaction ("carbonyl stress"). Carbonyl stress has been implicated in the content derived from the high glucose of the CAPD solution, from solutes being the end products of heat sterilization of the solution, and from uremic retention. Carbonyl stress might not only modify peritoneal matrix proteins and alter their structures, but also react with mesothelial and endothelial cell surface proteins and initiate a range of inflammatory responses. Carbonyl stress might therefore contribute to the development of peritoneal sclerosis in patients with long-term CAPD [reviewed in 13,14].

In CAPD patients, the peritoneal membrane is continuously exposed to the high glucose concentration contained in the dialysate, and this may lead to the local generation of AGEs. To test this hypothesis, the plasma and dialysate AGE concentrations in five CAPD patients

were evaluated. The dialysate was measured after a one-hour and after a 12-hour dwell time. Additionally, an immunohistochemical investigation of the peritoneal membrane for AGE was performed in two patients [15]. Only low concentrations of AGE were found in the dialysate after a one-hour dwell time; however, after 12 hours the dialysate AGE was even greater than the plasma concentration. Positive staining for AGE in the interstitium of the mesothelial layer was observed in both peritoneal specimens. The dialysate AGE contained a high proportion of high-molecular-weight AGE proteins and low-molecular-weight AGE was found to be in the same concentration range as the total serum AGE. It was therefore concluded that there is local generation of AGE in the peritoneal membrane and a "washing out" of AGE from the peritoneal membrane during longer dwell times. Other studies have also shown that formation of AGE products occurs in dialysis fluid and that factors such as pH and lactate in dialysate can modulate this process [16].

Another study confirmed that AGEs become predominantly accumulated in the vascular wall in accordance with the prolongation of CAPD treatment, and this might play a role in the increased permeability of the peritoneal membrane in CAPD [17].

In CAPD patients, AGE formation in the peritoneum correlates with the development of severe interstitial fibrosis and microvascular sclerosis, which is associated clinically with impaired peritoneal ultrafiltration. In patients with low peritoneal ultrafiltration, AGE accumulated in the peritoneal fibrous tissue and microvascular walls. Remarkably, AGE accumulated more intensely in hyalinized fibrosis of small venular media. The extent of AGE accumulation in peritoneal interstitium and vascular walls correlated with the progression of interstitial fibrosis and vascular sclerosis [18].

Furosine and pentosidine are examples of early and late glycation products, respectively. Plasma furosine and pentosidine were measured by HPLC in patients with renal dysfunction with or without diabetes mellitus, and dialysate pentosidine and furosine in CAPD patients. Plasma furosine was remarkably high in diabetes mellitus, hemodialysis, and CAPD patients. Plasma pentosidine appeared to depend on renal function and was not influenced by diabetic condition. Plasma pentosidine was significantly higher in CAPD than hemodialysis patients. A weak positive correlation was noted between dialysate and plasma furosine and pentosidine, indicating the main source of furosine and pentosidine in peritoneal dialysis effluent to be plasma. Thus, in situ formation of early glycation products in the peritoneal cavity takes place in CAPD patients, and high plasma pentosidine may lead to its accumulation in tissue, resulting, possibly, in pathological changes [19].

Pannekeet et al recently demonstrated that four-hour

pentosidine effluent contents are not influenced by dialysate glucose concentration or osmolality of the dialysate, in contrast to percentage glycation of albumin and IgG [20]. The relationship between the pentosidine effluent contents and duration of peritoneal dialysis, and the effect of nonglucose dialysate on the pentosidine effluent contents suggest that long-term glucose exposure is an important determinant of membrane glycosylation. Thus, pentosidine effluent contents probably reflect the long-term effects of intraperitoneal glycosylation of peritoneal membrane proteins. Six patients with peritoneal ultrafiltration failure were treated with nonglucose dialysis solutions and the pentosidine effluent contents were studied after six weeks. In five of six patients treated with nonglucose dialysate, pentosidine effluent contents decreased while serum pentosidine was stable, suggesting that treatment with nonglucose dialysis solutions may result in a "washout" of glycosylated proteins from the peritoneal membrane [20].

The peritoneal kinetics of pentosidine were first studied by Friedlander et al [21, 22]. The glycated content of peritoneal proteins was initially identical to plasma, but increased 200% within four hours due to in situ glycation, and this also was demonstrated in vitro. In contrast, peritoneal proteins contained a two to four times greater content of the AGE pentosidine at all equilibrium time points. It was concluded that the clearance of protein-associated pentosidine by the peritoneal membrane may explain the lower steady state levels in patients treated by peritoneal dialysis.

Miyata et al demonstrated that pentosidine accumulates markedly as an albumin-linked form and in a free-form in the plasma of patients with end-stage renal failure [23]. In a follow-up study, it appeared that hemodialysis cleared free pentosidine, but not albumin-linked pentosidine [24]. In contrast, CAPD cleared both forms. Plasma total pentosidine levels were significantly lower in CAPD than in hemodialysis, in part as the result of a lower serum albumin level in the former patients. However, the free pentosidine was also significantly lower in CAPD than in hemodialysis. The transport of both the free and albumin-linked pentosidine across the peritoneum occurs mainly by diffusion. Interestingly, peritoneal albumin-linked pentosidine clearance significantly exceeded albumin clearance. The fact that albumin-linked pentosidine levels were significantly higher in the peritoneal fluid than in the serum raises the intriguing possibility of a facilitated diffusion of albumin-linked pentosidine or an active transport mechanism for protein-linked pentosidine into the peritoneal cavity.

It has previously been shown that glucose solutions that were heat-sterilized showed an increase in UV absorbance at 284 nm and were cytotoxic [25]. These and subsequent results indicated that the toxic products formed during heat sterilization of PD fluids are derived

from glucose [25, 26]. Among these toxic products, the dicarbonyl compound 3-DG, is produced. The 3-DG concentrations in commercially and laboratory-prepared heat-sterilized fluids were 118 and 154  $\mu\text{mol/L}$ , respectively. Gambrosol-Bio dialysate produced through an alteration of the manufacturing condition, and sterile-filtered fluid produced in the laboratory contained 3-DG in concentrations of 12.3 and less than 1.2  $\mu\text{mol/L}$ , respectively. It can be speculated that the presence of 3-DG in unused conventional peritoneal dialysis-fluid could act as a local promoter, and increase local AGE formation within the peritoneal cavity [27].

In vitro studies have shown that heat-sterilized icodextrin (polyglucose) peritoneal dialysis fluid results in less glycation and AGE formation than conventional heat-sterilized glucose dialysate [28].

### ADVANCED OXIDATION PROTEIN PRODUCTS


Like lipids, amino acids, peptides and proteins are vulnerable to attacks by a variety of free radicals and related oxidants, and high levels of advanced oxidation protein products have been found in the plasma of uremic patients [29–31]. Interestingly, the plasma levels of these AOPPs were lower in CAPD patients than in hemodialysis patients. Furthermore, a close correlation was found between the AOPP levels and the AGE-pentosidine levels linking the Maillard reaction to oxidative pathway of injury to biological molecules [29].

### GRANULOCYTE INHIBITORY PROTEINS

A peptide that inhibits the chemotactic movement of polymorphonuclear leukocytes in a concentration dependent, nonreversible manner was recently isolated from the peritoneal effluent of CAPD patients [32, 33]. Amino acid sequencing showed that the isolated peptide has the same amino-terminal sequence as ubiquitin. The same group had previously isolated two other granulocyte inhibitory proteins (GIP I and GIP II) with homology to light chain proteins and  $\beta_2\text{m}$ , respectively [34]. It was speculated that these proteins were responsible, at least in part, for impaired local cellular host defense in the peritoneal cavity.

### p-CRESOL

p-Cresol is a phenolic and volatile compound with a molecular weight of only 108.1 D. It is partially lipophilic and binds strongly to plasma proteins (close to 100%) under normal conditions. p-Cresol is an end product of protein breakdown produced by the intestinal bacteria, as a result of the metabolism of tyrosine and phenylalanine, and an increase of the nutritional protein load in



**Fig. 1. Plasma levels of total (T) and free (F) p-Cresol in continuous ambulatory peritoneal dialysis (CAPD) and hemodialysis (HD) patients. (A)** p-Cresol levels in two groups of CAPD patients with different residual renal function. **(B)** p-Cresol levels in CAPD and hemodialysis patients matched for residual renal function. The plasma levels of total p-Cresol are significantly higher in the hemodialysis patients. **(C)** p-Cresol plasma levels in CAPD patients and anuric patients treated with hemodialysis. Both the total and free levels of p-Cresol are significantly higher in the hemodialysis patients.

healthy individuals results in its enhanced generation and urinary excretion [35]. The serum concentration of p-Cresol is elevated in renal failure [2]. Recently, a sensitive method to determine total and free p-Cresol has been developed [36]. As illustrated in Figure 1, the total and free levels of p-Cresol seemed to be lower in CAPD compared to levels found in hemodialysis. These differences remained significant even after correction for differences in residual renal function between peritoneal dialysis and hemodialysis patients. At present, p-Cresol has been linked to several disturbances of biological

functions in uremia, amongst others the phagocytic response to bacterial load [reviewed in 35].

## HYPERHOMOCYSTEINEMIA

The high prevalence of cardiovascular disease in ESRD that is not explained by the classic risk factors, has been clarified considerably by the linkage of hyperhomocysteinemia, a common finding in ESRD, with an increased risk of cardiac disease and atherosclerosis [37–39].

Total homocysteine (Hcy) values averaged 29.8  $\mu\text{mol/L}$  in hemodialysis patients, significantly higher than the mean value of 19.9  $\mu\text{mol/L}$  observed in patients on peritoneal dialysis ( $P < 0.001$ ). The prevalence of hyperhomocysteinemia was 90.8% among hemodialysis patients, significantly higher than the prevalence of 67.4% among peritoneal dialysis patients. Folate values in hemodialysis patients averaged 45.5 nmol/L and were significantly lower than in peritoneal dialysis patients [40]. Folate deficiency has been linked to hyperhomocysteinemia.

A common polymorphism in the 5,10-methylenetetrahydrofolate reductase (*MTHFR*) gene (C677T transition) results in increased total homocysteine levels in peritoneal dialysis patients compared to age- and sex-matched healthy individuals. The allelic frequency of the C677T transition in the *MTHFR* gene in peritoneal dialysis patients (0.29) was, however, comparable to the frequency in healthy individuals (0.34) [41].

The peritoneal elimination of total homocysteine primarily depends on the plasma total homocysteine concentration. Elevated total homocysteine plasma levels cannot be reduced efficiently by peritoneal dialysis [42]. In contrast, a significant and prolonged reduction in Hcy concentrations by peritoneal dialysis in ESRD patients was observed by Ducloux et al [43]. The decrease in Hcy concentration was positively related to dialysis adequacy. This study suggests the possibility that dialysis adequacy may influence arteriosclerotic outcomes through a Hcy-lowering effect.

Hyperhomocysteinemia can be normalized with folic acid alone in about 40% of peritoneal dialysis patients. Betaine does not further lower plasma homocysteine levels. A maintenance dose of 1 or 5 mg folic acid daily results in equivalent plasma homocysteine levels. Long-term reduction in plasma homocysteine did not result in an improvement of endothelial function as assessed by determination of endothelium-dependent vasodilation and biochemical markers [44].

Methionine-containing amino acid containing peritoneal fluid induces an increase in the plasma total homocysteine level. This has the potential to offset the beneficial effects of an improved serum lipid profile and reduced fat mass on the risk of cardiovascular disease in peritoneal dialysis patients. Therefore, lowering the methionine con-

tent of the fluid may be required to overcome this adverse effect [45].

## ROLE OF UREMIC TOXINS IN THE PROGRESSION OF RENAL FUNCTION

An interesting study explored the possibility that circulating uremic toxins were involved in the progressive loss of intact nephrons in chronic renal failure [46]. This loss is attributed to progressive sclerosis at the level of the glomeruli of the intact nephrons. Rats underwent subtotal nephrectomy and a kidney biopsy after seven weeks and the extent of glomerular sclerosis was semiquantitatively estimated. One group of animals was treated with peritoneal dialysis, while another group received an oral charcoal adsorbent, called AST-120. The animals were then re-examined by micropuncture and histology at week 12. The group of animals that was treated with peritoneal dialysis showed lesser glomerulosclerosis than the control group at the end of the experiment and the percentage of glomeruli with a higher sclerosis index was much higher in the undialyzed animals. Also, the glomerular filtration rate was significantly higher at 12 weeks in the dialyzed animals. These results were taken as evidence that biologically active circulating substances released in uremia are involved in the progression of glomerular sclerosis and that peritoneal dialysis can at least partially remove these substances.

Other uremic toxins like albumin-bound furancarboxylic acid, an inhibitor of erythropoiesis, show lower blood levels in peritoneal dialysis compared to hemodialysis patients, and this can also be explained by a more efficient removal across the peritoneal membrane [47].

Reprint requests to Norbert Lameire, M.D., Renal Division, Department of Medicine, University Hospital, 185, De Pintelaan, 9000 Gent, Belgium.

## REFERENCES

- BERGSTRÖM J: Uremic toxicity. Nutritional Management of Renal Disease, edited by KOPPLE JD, MASSRY SG, Baltimore, Williams & Wilkins, p 97, 1997
- VANHOLDER R: Uremic toxins. *Adv Nephrol Necker Hosp* 26:143-163, 1997
- VANHOLDER R, DE SMET R: Pathophysiologic effects of uremic retention solutes. *J Am Soc Nephrol* 10:1815-1823, 1999
- LAMEIRE NH: The impact of residual renal function on the adequacy of peritoneal dialysis. *Nephron* 77:13-28, 1997
- KABANDA A, GOFFIN E, BERNARD A, LAUWERYS R, VAN YPERSELE DE STRIHOUC: Factors influencing serum levels and peritoneal clearances of low molecular weight proteins in continuous ambulatory peritoneal dialysis. *Kidney Int* 48:1946-1952, 1995
- CATIZONE L, VIGLIETTA G, COCCHI R, DADONE C, DE VA GIAN-GRANDE A, et al: Dialysis-related amyloidosis in a large CAPD population. *Adv Perit Dial* 11:213-217, 1995
- DIRAIMONDO CR, MCCARLEY P, STONE WJ: Beta-2 microglobulin in peritoneal dialysis patients: Serum levels and peritoneal clearances. *Perit Dial Int* 8:43-47, 1988
- WADA T, MIYATA T, SAKAI H, KUROKAWA K: Beta2-microglobulin and renal bone disease. *Perit Dial Int* 19 (Suppl 2):S413-S416, 1999
- NOMOTO Y, KAWAGUCHI Y, OHIRA S, YURI T, KUBO H, KUBOTA M, et al: Carpal tunnel syndrome in patients undergoing CAPD: A collaborative study in 143 centers. *Am J Nephrol* 15:295-299, 1995
- JADOUL M, GARBAR C, VANHOLDER R, SENNESAEEL J, MICHEL C, ROBERT A, et al: Prevalence of histological beta2-microglobulin amyloidosis in CAPD patients compared with hemodialysis patients. *Kidney Int* 54:956-959, 1998
- TAN SY, BAILLOD R, BROWN E, FARRINGTON K, SOPER C, PERCY M, et al: Clinical, radiological and serum amyloid P component scintigraphic features of beta2-microglobulin amyloidosis associated with continuous ambulatory peritoneal dialysis. *Nephrol Dial Transplant* 14:1467-1471, 1999
- NIWA T, KATSUZAKI T, MOMOI T, MIYAZAKI T, OGAWA H, SAITO A, et al: Modification of beta 2m with advanced glycation end products as observed in dialysis-related amyloidosis by 3-DG accumulating in uremic serum. *Kidney Int* 49:861-867, 1996
- MIYATA T, IZUHARA Y, SAKAI H, KUROKAWA K: Carbonyl stress: Increased carbonyl modification of tissue and cellular proteins in uremia. *Perit Dial Int* 19(Suppl 2):S58-S61, 1999
- DAWNAY A, MILLAR DJ: The pathogenesis and consequences of AGE formation in uraemia and its treatment. *Cell Mol Biol* 44:1081-1094, 1998
- MAHIOUT A, EHLENDING G, BRUNKHORST R: Advanced glycation end-products in the peritoneal fluid and in the peritoneal membrane of continuous ambulatory peritoneal dialysis patients. *Nephrol Dial Transplant* 11(Suppl 5):2-6, 1996
- LAMB EJ, CATTELL WR, DAWNAY AB: In vitro formation of advanced glycation end products in peritoneal dialysis fluid. *Kidney Int* 47:1768-1774, 1995
- NAKAYAMA M, KAWAGUCHI Y, YAMADA K, HASEGAWA T, TAKAZOE K, KATOH N et al: Immunohistochemical detection of advanced glycosylation end-products in the peritoneum and its possible pathophysiological role in CAPD. *Kidney Int* 51:182-186, 1997
- HONDA K, NITTA K, HORITA S, YUMURA W, NIHEI H, NAGAI R et al: Accumulation of advanced glycation end products in the peritoneal vasculature of continuous ambulatory peritoneal dialysis patients with low ultra-filtration. *Nephrol Dial Transplant* 14:1541-1549, 1999
- KUMANO K, YOKOTA S, SAKAI T, KOBAYASHI N, YOSHIDA A, YOSHIMURA T, et al: Kinetic analysis of furosine and pentosidine in CAPD patients. *Adv Perit Dial* 13:53-57, 1997
- HO DAC PANNEKEET M, WEISS MF, DE WAART DR, ERHARD P, HIRALALL JK, KREDIET RT: Analysis of non enzymatic glycosylation in vivo: Impact of different dialysis solutions. *Perit Dial Int* 19(Suppl 2):S68-S741, 1999
- FRIEDLANDER MA, WU YC, SCHULAK YA, MONNIER VM, HRICIK DA: Influence of dialysis modality on plasma and tissue concentrations of pentosidine in patients with end-stage renal disease. *Am J Kidney Dis* 25:441-451, 1995
- FRIEDLANDER MA, WU YC, ELGAWISH A, MONNIER VM: Early and advanced glycosylation end products. Kinetics of formation and clearance in peritoneal dialysis. *J Clin Invest* 97:728-735, 1996
- MIYATA T, UEDA Y, SHINZATO T, IIDA Y, TANAKA S, KUROKAWA K, et al: Accumulation of albumin-linked and free-form pentosidine in the circulation of uremic patients with end-stage renal failure: Renal implications in the pathophysiology of pentosidine. *J Am Soc Nephrol* 7:1198-1206, 1996
- MIYATA T, UEDA Y, YOSHIDA A, SUGIYAMA S, IIDA Y, JADOUL M, et al: Clearance of pentosidine, an advanced glycation end product, by different modalities of renal replacement therapy. *Kidney Int* 51:880-887, 1997
- MARTINSON E, WIESLANDER A, KJELLSTRAND P, BOBERG U: Toxicity of heat sterilized peritoneal dialysis fluids is derived from degradation of glucose. *ASAIO J* 38:M370-M372, 1992
- WIESLANDER AP: Cytotoxicity of peritoneal dialysis fluid—is it related to glucose breakdown products? [editorial] *Nephrol Dial Transplant* 11:958-959, 1996
- LINDEN T, FORSBÄCK G, DEPPISCH R, HENLE T, WIESLANDER A: 3-Deoxyglucosone, a promoter of advanced glycation end products in fluids for peritoneal dialysis. *Perit Dial Int* 18:290-293, 1998
- DAWNAY AB, MILLAR DJ: Glycation and advanced glycation end-product formation with icodextrin and dextrose. *Perit Dial Int* 17:52-58, 1997
- WITKO-SARSAT V, FRIEDLANDER M, CAPELLÈRE-BLANDIN C, NGU-

- YEN-KHOA T, NGUYEN AT, ZINGRAFF J, *et al*: Advanced oxidation protein products as a novel marker of oxidative stress in uremia. *Kidney Int* 49:1304-1313, 1996
30. WITKO-SARSAT V, NGUYEN-KHOA T, JUNGERS P, DRUEKE TB, DESCAMPS-LATSCHA B: Advanced oxidation protein products as a novel molecular basis of oxidative stress in uraemia. *Nephrol Dial Transplant* 14(Suppl 1):76-78, 1999
  31. WITKO-SARSAT V, NGUYEN-KHOA T, NGUYEN AT, JUNGERS P, DRUEKE T, DESCAMPS-LATSCHA B: Advanced oxidation protein products as a novel marker of oxidative stress: Oxidative stress markers and mediators of inflammation in uremia. *Adv Nephrol Necker Hosp* 28:321-341, 1998
  32. COHEN G, RUDNICKI M, HÖRL WH: A peptide isolated from a patient on continuous ambulatory peritoneal dialysis has homology to ubiquitin and inhibits the chemotactic response of polymorphonuclear leukocytes. *Miner Electrolyte Metab* 23:210-213, 1997
  33. COHEN G, RUDNICKI M, HÖRL WH: Isolation of modified ubiquitin as a neutrophil chemotaxis inhibitor from uremic patients. *J Am Soc Nephrol* 9:451-456, 1998
  34. HAAG-WEBER M, MAJ B, HÖRL WH: Impaired cellular host defense in peritoneal dialysis by two granulocyte inhibitory proteins. *Nephrol Dial Transplant* 9:1769-1773, 1994
  35. VANHOLDER R, DE SMET R, LESAFFER G: p-Cresol: A toxin revealing many neglected but relevant aspects of uraemic toxicity. *Nephrol Dial Transplant* 14:2813-2815, 1999
  36. DE SMET R, GLORIEUX G, HSU C, VANHOLDER R: p-Cresol and uric acid: Two old uremic toxins revisited. *Kidney Int* 53(Suppl 62):S8-S11, 1997
  37. GUPTA A, ROBINSON K: Hyperhomocysteinaemia and end stage renal disease. *J Nephrol* 10:77-84, 1997
  38. KRONENBERG F: Homocysteine, lipoprotein (a) and fibrinogen: Metabolic risk factors for cardiovascular complications of chronic renal disease. *Curr Opin Nephrol Hypertens* 7:271-278, 1998
  39. BOSTOM AG, LATHROP L: Hyperhomocysteinemia in end-stage renal disease: Prevalence, etiology, and potential relationship to atherosclerotic outcomes. *Kidney Int* 52:10-20, 1997
  40. MOUSTAPHA A, GUPTA A, ROBINSON K, ARHEART K, JACOBSEN DW, SCHREIBER MJ, *et al*: Prevalence and determinants of hyperhomocysteinemia in hemodialysis and peritoneal dialysis. *Kidney Int* 55:1470-1475, 1999
  41. VYCHYTIL A, FÖDINGER M, WÖFL G, ENZENBERGER B, AUINGER M, PRISCHL F, *et al*: Major determinants of hyperhomocysteinemia in peritoneal dialysis patients. *Kidney Int* 53:1775-1782, 1998
  42. VYCHYTIL A, FÖDINGER M, PAPAGIANNOPOULOS M, WÖFL G, HÖRL WH, SUNDER-PLESSMANN G: Peritoneal elimination of homocysteine moieties in continuous ambulatory peritoneal dialysis patients. *Kidney Int* 55:2054-2061, 1999
  43. DUCLOUX D, HEUZE-LECORNU L, GIBEY R, BRESSON-VAUTRIN C, VAUTRIN P, CHALOPIN JM: Dialysis adequacy and homocyst(e)ine concentrations in peritoneal dialysis patients. *Nephrol Dial Transplant* 14:728-731, 1999
  44. VAN GULDENER C, JANSSEN MJ, LAMBERT J, TER WEE P, DONKER AJ, STEHOUWER CD: Folic acid treatment of hyperhomocysteinemia in peritoneal dialysis patients: No change in endothelial function after long-term therapy. *Perit Dial Int* 18:282-289, 1998
  45. BRULEZ HF, VAN GULDENER C, DONKER AJ, TER WEE P: The impact of an amino acid-based peritoneal dialysis fluid on plasma total homocysteine levels, lipid profile and body fat mass. *Nephrol Dial Transplant* 14:154-159, 1999
  46. MOTOJIMA M, NISHIJIMA F, IKOMA M, KAWAMURA T, YOSHIOKA T, FOGO AB, *et al*: Role for 'uremic toxin' in the progressive loss of intact nephrons in chronic renal failure. *Kidney Int* 40:461-469, 1991
  47. NIWA T, YAZAWA T, KODAMA T, UEHARA Y, MAEDA K, YAMADA K: Efficient removal of albumin-bound furancarboxylic acid, an inhibitor of erythropoiesis, by continuous ambulatory peritoneal dialysis. *Nephron* 56:241-245, 1990